

Finding Relationships Among Variables

BUS 230: Business Research and Communication

- Specific goals:
 - Re-familiarize ourselves with basic statistics ideas: sampling distributions, hypothesis tests, p-values.
 - Be able to distinguish different types of data and prescribe appropriate statistical methods.
 - Conduct a number of hypothesis tests using methods appropriate for questions involving only one or two variables.
- Learning objectives:
 - LO2: Interpret data using statistical analysis.
 - LO2.3: Formulate conclusions and recommendations based upon statistical results.

- Specific goals:
 - Re-familiarize ourselves with basic statistics ideas: sampling distributions, hypothesis tests, p-values.
 - Be able to distinguish different types of data and prescribe appropriate statistical methods.
 - Conduct a number of hypothesis tests using methods appropriate for questions involving only one or two variables.
- Learning objectives:
 - LO2: Interpret data using statistical analysis.
 - LO2.3: Formulate conclusions and recommendations based upon statistical results.

What to Look For

- For each test, remember the following:
 - In plain English, be able to describe the purpose of the test.
 - Know whether the test is a parametric test or a non-parametric test.
 - Know the null and alternative hypotheses.
- When choosing a test to answer a research question, ask yourself:
 - ① What is your research question?
 - ② How many variables do you have?
 - ③ What is your scale of measurement?
 - ④ Are you looking for differences or other relationship?
 - ⑤ If you are looking for differences, are your observations independent or paired?
 - ⑥ Do the hypotheses (or the result of the hypothesis test) answer your research question?

What to Look For

- For each test, remember the following:
 - In plain English, be able to describe the purpose of the test.
 - Know whether the test is a parametric test or a non-parametric test.
 - Know the null and alternative hypotheses.
- When choosing a test to answer a research question, ask yourself:
 - ① What is your research question?
 - ② How many variables do you have?
 - ③ What is your scale of measurement?
 - ④ Are you looking for differences or other relationship?
 - ⑤ If you are looking for differences, are your observations independent or paired?
 - ⑥ Do the hypotheses (or the result of the hypothesis test) answer your research question?

What to Look For

- For each test, remember the following:
 - In plain English, be able to describe the purpose of the test.
 - Know whether the test is a parametric test or a non-parametric test.
 - Know the null and alternative hypotheses.
- When choosing a test to answer a research question, ask yourself:
 - ① What is your research question?
 - ② How many variables do you have?
 - ③ What is your scale of measurement?
 - ④ Are you looking for differences or other relationship?
 - ⑤ If you are looking for differences, are your observations independent or paired?
 - ⑥ Do the hypotheses (or the result of the hypothesis test) answer your research question?

What to Look For

- For each test, remember the following:
 - In plain English, be able to describe the purpose of the test.
 - Know whether the test is a parametric test or a non-parametric test.
 - Know the null and alternative hypotheses.
- When choosing a test to answer a research question, ask yourself:
 - ① What is your research question?
 - ② How many variables do you have?
 - ③ What is your scale of measurement?
 - ④ Are you looking for differences or other relationship?
 - ⑤ If you are looking for differences, are your observations independent or paired?
 - ⑥ Do the hypotheses (or the result of the hypothesis test) answer your research question?

What to Look For

- For each test, remember the following:
 - In plain English, be able to describe the purpose of the test.
 - Know whether the test is a parametric test or a non-parametric test.
 - Know the null and alternative hypotheses.
- When choosing a test to answer a research question, ask yourself:
 - 1 What is your research question?
 - 2 How many variables do you have?
 - 3 What is your scale of measurement?
 - 4 Are you looking for differences or other relationship?
 - 5 If you are looking for differences, are your observations independent or paired?
 - 6 Do the hypotheses (or the result of the hypothesis test) answer your research question?

What to Look For

- For each test, remember the following:
 - In plain English, be able to describe the purpose of the test.
 - Know whether the test is a parametric test or a non-parametric test.
 - Know the null and alternative hypotheses.
- When choosing a test to answer a research question, ask yourself:
 - 1 What is your research question?
 - 2 How many variables do you have?
 - 3 What is your scale of measurement?
 - 4 Are you looking for differences or other relationship?
 - 5 If you are looking for differences, are your observations independent or paired?
 - 6 Do the hypotheses (or the result of the hypothesis test) answer your research question?

What to Look For

- For each test, remember the following:
 - In plain English, be able to describe the purpose of the test.
 - Know whether the test is a parametric test or a non-parametric test.
 - Know the null and alternative hypotheses.
- When choosing a test to answer a research question, ask yourself:
 - 1 What is your research question?
 - 2 How many variables do you have?
 - 3 What is your scale of measurement?
 - 4 Are you looking for differences or other relationship?
 - 5 If you are looking for differences, are your observations independent or paired?
 - 6 Do the hypotheses (or the result of the hypothesis test) answer your research question?

What to Look For

- For each test, remember the following:
 - In plain English, be able to describe the purpose of the test.
 - Know whether the test is a parametric test or a non-parametric test.
 - Know the null and alternative hypotheses.
- When choosing a test to answer a research question, ask yourself:
 - 1 What is your research question?
 - 2 How many variables do you have?
 - 3 What is your scale of measurement?
 - 4 Are you looking for differences or other relationship?
 - 5 If you are looking for differences, are your observations independent or paired?
 - 6 Do the hypotheses (or the result of the hypothesis test) answer your research question?

What to Look For

- For each test, remember the following:
 - In plain English, be able to describe the purpose of the test.
 - Know whether the test is a parametric test or a non-parametric test.
 - Know the null and alternative hypotheses.
- When choosing a test to answer a research question, ask yourself:
 - 1 What is your research question?
 - 2 How many variables do you have?
 - 3 What is your scale of measurement?
 - 4 Are you looking for differences or other relationship?
 - 5 If you are looking for differences, are your observations independent or paired?
 - 6 Do the hypotheses (or the result of the hypothesis test) answer your research question?

What to Look For

- For each test, remember the following:
 - In plain English, be able to describe the purpose of the test.
 - Know whether the test is a parametric test or a non-parametric test.
 - Know the null and alternative hypotheses.
- When choosing a test to answer a research question, ask yourself:
 - 1 What is your research question?
 - 2 How many variables do you have?
 - 3 What is your scale of measurement?
 - 4 Are you looking for differences or other relationship?
 - 5 If you are looking for differences, are your observations independent or paired?
 - 6 Do the hypotheses (or the result of the hypothesis test) answer your research question?

What to Look For

- For each test, remember the following:
 - In plain English, be able to describe the purpose of the test.
 - Know whether the test is a parametric test or a non-parametric test.
 - Know the null and alternative hypotheses.
- When choosing a test to answer a research question, ask yourself:
 - 1 What is your research question?
 - 2 How many variables do you have?
 - 3 What is your scale of measurement?
 - 4 Are you looking for differences or other relationship?
 - 5 If you are looking for differences, are your observations independent or paired?
 - 6 Do the hypotheses (or the result of the hypothesis test) answer your research question?

Correlation

- A **correlation** exists between two variables when one of them is related to the other in some way, such that there is **co-movement**.
- The **Pearson linear correlation coefficient** is a measure of the strength of the linear relationship between two variables.
 - Parametric test!
 - Null hypothesis: there is zero linear correlation between two variables.
 - Alternative hypothesis: there is a linear correlation (either positive or negative) between two variables.
- Spearman's Rank Test
 - Non-parametric test.
 - Behind the scenes - replaces actual data with their *rank*, computes the Pearson using ranks.
 - Same hypotheses.

Correlation

- A **correlation** exists between two variables when one of them is related to the other in some way, such that there is **co-movement**.
- The **Pearson linear correlation coefficient** is a measure of the strength of the linear relationship between two variables.
 - Parametric test!
 - Null hypothesis: there is zero linear correlation between two variables.
 - Alternative hypothesis: there is a linear correlation (either positive or negative) between two variables.
- Spearman's Rank Test
 - Non-parametric test.
 - Behind the scenes - replaces actual data with their *rank*, computes the Pearson using ranks.
 - Same hypotheses.

Correlation

- A **correlation** exists between two variables when one of them is related to the other in some way, such that there is **co-movement**.
- The **Pearson linear correlation coefficient** is a measure of the strength of the linear relationship between two variables.
 - Parametric test!
 - Null hypothesis: there is zero linear correlation between two variables.
 - Alternative hypothesis: there is a linear correlation (either positive or negative) between two variables.
- Spearman's Rank Test
 - Non-parametric test.
 - Behind the scenes - replaces actual data with their *rank*, computes the Pearson using ranks.
 - Same hypotheses.

Correlation

- A **correlation** exists between two variables when one of them is related to the other in some way, such that there is **co-movement**.
- The **Pearson linear correlation coefficient** is a measure of the strength of the linear relationship between two variables.
 - Parametric test!
 - Null hypothesis: there is zero linear correlation between two variables.
 - Alternative hypothesis: there is a linear correlation (either positive or negative) between two variables.
- Spearman's Rank Test
 - Non-parametric test.
 - Behind the scenes - replaces actual data with their *rank*, computes the Pearson using ranks.
 - Same hypotheses.

Correlation

- A **correlation** exists between two variables when one of them is related to the other in some way, such that there is **co-movement**.
- The **Pearson linear correlation coefficient** is a measure of the strength of the linear relationship between two variables.
 - Parametric test!
 - Null hypothesis: there is zero linear correlation between two variables.
 - Alternative hypothesis: there is a linear correlation (either positive or negative) between two variables.
- Spearman's Rank Test
 - Non-parametric test.
 - Behind the scenes - replaces actual data with their *rank*, computes the Pearson using ranks.
 - Same hypotheses.

Correlation

- A **correlation** exists between two variables when one of them is related to the other in some way, such that there is **co-movement**.
- The **Pearson linear correlation coefficient** is a measure of the strength of the linear relationship between two variables.
 - Parametric test!
 - Null hypothesis: there is zero linear correlation between two variables.
 - Alternative hypothesis: there is a linear correlation (either positive or negative) between two variables.
- Spearman's Rank Test
 - Non-parametric test.
 - Behind the scenes - replaces actual data with their *rank*, computes the Pearson using ranks.
 - Same hypotheses.

Correlation

- A **correlation** exists between two variables when one of them is related to the other in some way, such that there is **co-movement**.
- The **Pearson linear correlation coefficient** is a measure of the strength of the linear relationship between two variables.
 - Parametric test!
 - Null hypothesis: there is zero linear correlation between two variables.
 - Alternative hypothesis: there is a linear correlation (either positive or negative) between two variables.
- Spearman's Rank Test
 - Non-parametric test.
 - Behind the scenes - replaces actual data with their *rank*, computes the Pearson using ranks.
 - Same hypotheses.


Correlation

- A **correlation** exists between two variables when one of them is related to the other in some way, such that there is **co-movement**.
- The **Pearson linear correlation coefficient** is a measure of the strength of the linear relationship between two variables.
 - Parametric test!
 - Null hypothesis: there is zero linear correlation between two variables.
 - Alternative hypothesis: there is a linear correlation (either positive or negative) between two variables.
- Spearman's Rank Test
 - Non-parametric test.
 - Behind the scenes - replaces actual data with their *rank*, computes the Pearson using ranks.
 - Same hypotheses.


Correlation

- A **correlation** exists between two variables when one of them is related to the other in some way, such that there is **co-movement**.
- The **Pearson linear correlation coefficient** is a measure of the strength of the linear relationship between two variables.
 - Parametric test!
 - Null hypothesis: there is zero linear correlation between two variables.
 - Alternative hypothesis: there is a linear correlation (either positive or negative) between two variables.
- Spearman's Rank Test
 - Non-parametric test.
 - Behind the scenes - replaces actual data with their *rank*, computes the Pearson using ranks.
 - Same hypotheses.


Positive linear correlation



(a) Positive correlation between x and y




(b) Strong positive correlation between x and y




(c) Perfect positive correlation between x and y

- Positive correlation: two variables move in the same direction.
- Stronger the correlation: closer the correlation coefficient is to 1.
- Perfect positive correlation: $\rho = 1$


Positive linear correlation



(a) Positive correlation between x and y




(b) Strong positive correlation between x and y




(c) Perfect positive correlation between x and y

- Positive correlation: two variables move in the same direction.
- Stronger the correlation: closer the correlation coefficient is to 1.
- Perfect positive correlation: $\rho = 1$


Positive linear correlation



(a) Positive correlation between x and y




(b) Strong positive correlation between x and y




(c) Perfect positive correlation between x and y

- Positive correlation: two variables move in the same direction.
- Stronger the correlation: closer the correlation coefficient is to 1.
- Perfect positive correlation: $\rho = 1$


Negative linear correlation



(d) Negative correlation between x and y




(e) Strong negative correlation between x and y




(f) Perfect negative correlation between x and y

- Negative correlation: two variables move in opposite directions.
- Stronger the correlation: closer the correlation coefficient is to -1 .
- Perfect negative correlation: $\rho = -1$


Negative linear correlation



(d) Negative correlation between x and y




(e) Strong negative correlation between x and y




(f) Perfect negative correlation between x and y

- Negative correlation: two variables move in opposite directions.
- Stronger the correlation: closer the correlation coefficient is to -1 .
- Perfect negative correlation: $\rho = -1$


Negative linear correlation



(d) Negative correlation between x and y




(e) Strong negative correlation between x and y




(f) Perfect negative correlation between x and y

- Negative correlation: two variables move in opposite directions.
- Stronger the correlation: closer the correlation coefficient is to -1 .
- Perfect negative correlation: $\rho = -1$

No linear correlation




(g) No correlation between x and y




(h) Nonlinear relationship between x and y

- Panel (g): no relationship at all.
- Panel (h): strong relationship, but not a *linear* relationship.
 - Cannot use regular correlation to detect this.

No linear correlation



(g) No correlation between x and y



(h) Nonlinear relationship between x and y

- Panel (g): no relationship at all.
- Panel (h): strong relationship, but not a *linear* relationship.
 - Cannot use regular correlation to detect this.

Chi-Squared Test for Independence

- Used to determine if two categorical variables (eg: nominal) are related.
- Example: Suppose a hotel manager surveys guest who indicate they will not return:

Reason for Stay	Reason for Not Returning		
	Price	Location	Amenities
Personal/Vacation	56	49	0
Business	20	47	27

- Data in the table are always frequencies that fall into individual categories.
- Could use this table to test if two variables are independent.

Chi-Squared Test for Independence

- Used to determine if two categorical variables (eg: nominal) are related.
- Example: Suppose a hotel manager surveys guest who indicate they will not return:

Reason for Stay	Reason for Not Returning		
	Price	Location	Amenities
Personal/Vacation	56	49	0
Business	20	47	27

- Data in the table are always frequencies that fall into individual categories.
- Could use this table to test if two variables are independent.

Chi-Squared Test for Independence

- Used to determine if two categorical variables (eg: nominal) are related.
- Example: Suppose a hotel manager surveys guest who indicate they will not return:

Reason for Not Returning

Reason for Stay	Price	Location	Amenities
Personal/Vacation	56	49	0
Business	20	47	27

- Data in the table are always frequencies that fall into individual categories.
- Could use this table to test if two variables are independent.

Chi-Squared Test for Independence

- Used to determine if two categorical variables (eg: nominal) are related.
- Example: Suppose a hotel manager surveys guest who indicate they will not return:

Reason for Not Returning

Reason for Stay	Price	Location	Amenities
Personal/Vacation	56	49	0
Business	20	47	27

- Data in the table are always frequencies that fall into individual categories.
- Could use this table to test if two variables are independent.

Chi-Squared Test for Independence

- Used to determine if two categorical variables (eg: nominal) are related.
- Example: Suppose a hotel manager surveys guest who indicate they will not return:

Reason for Not Returning

Reason for Stay	Price	Location	Amenities
Personal/Vacation	56	49	0
Business	20	47	27

- Data in the table are always frequencies that fall into individual categories.
- Could use this table to test if two variables are independent.

Test of independence

- **Null hypothesis:** there is no relationship between the row variable and the column variable (independent)
- **Alternative hypothesis:** There is a relationship between the row variable and the column variable (dependent).
- Test statistic:

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

- O : observed frequency in a cell from the contingency table.
- E : expected frequency computed with the *assumption that the variables are independent*.
- Large χ^2 values indicate variables are dependent (reject the null hypothesis).

Test of independence

- **Null hypothesis:** there is no relationship between the row variable and the column variable (independent)
- **Alternative hypothesis:** There is a relationship between the row variable and the column variable (dependent).
- Test statistic:

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

- O : observed frequency in a cell from the contingency table.
- E : expected frequency computed with the *assumption that the variables are independent*.
- Large χ^2 values indicate variables are dependent (reject the null hypothesis).

Test of independence

- **Null hypothesis:** there is no relationship between the row variable and the column variable (independent)
- **Alternative hypothesis:** There is a relationship between the row variable and the column variable (dependent).
- Test statistic:

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

- O : observed frequency in a cell from the contingency table.
- E : expected frequency computed with the *assumption that the variables are independent*.
- Large χ^2 values indicate variables are dependent (reject the null hypothesis).

Test of independence

- **Null hypothesis:** there is no relationship between the row variable and the column variable (independent)
- **Alternative hypothesis:** There is a relationship between the row variable and the column variable (dependent).
- Test statistic:

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

- O : observed frequency in a cell from the contingency table.
- E : expected frequency computed with the *assumption that the variables are independent*.
- Large χ^2 values indicate variables are dependent (reject the null hypothesis).

Test of independence

- **Null hypothesis:** there is no relationship between the row variable and the column variable (independent)
- **Alternative hypothesis:** There is a relationship between the row variable and the column variable (dependent).
- Test statistic:

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

- O : observed frequency in a cell from the contingency table.
- E : expected frequency computed with the *assumption that the variables are independent*.
- Large χ^2 values indicate variables are dependent (reject the null hypothesis).

Test of independence

- **Null hypothesis:** there is no relationship between the row variable and the column variable (independent)
- **Alternative hypothesis:** There is a relationship between the row variable and the column variable (dependent).
- Test statistic:

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

- O : observed frequency in a cell from the contingency table.
- E : expected frequency computed with the *assumption that the variables are independent*.
- Large χ^2 values indicate variables are dependent (reject the null hypothesis).

Test of independence

- **Null hypothesis:** there is no relationship between the row variable and the column variable (independent)
- **Alternative hypothesis:** There is a relationship between the row variable and the column variable (dependent).
- Test statistic:

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

- O : observed frequency in a cell from the contingency table.
- E : expected frequency computed with the *assumption that the variables are independent*.
- Large χ^2 values indicate variables are dependent (reject the null hypothesis).