

Overview of Statistical Methods / ANOVA

BUS 735: Business Decision Making and Research

1

1.1 Goals

Goals

- Specific goals:
 - Re-familiarize ourselves with statistical tests.
 - Learn how to choose appropriate tests.
 - Learn how to compare means or medians among more than two populations.
- Learning objectives:
 - LO1: Be able to construct and test hypotheses using a variety of bivariate statistical methods to compare characteristics between two populations.
 - LO3: Be able to construct and use analysis of variance and analysis of covariance models to construct and test hypotheses considering complex relationships among multiple variables.
 - LO6: Be able to use standard computer packages such as SPSS and Excel to conduct the quantitative analyses described in the learning objectives above.

2 Selecting the Right Method

Selecting Right Method

- Parametric Methods:
 - Only for *interval or ratio data*.
 - Make sure assumptions of CLT hold:
 - * Large sample size *or*.
 - * Normal distributed *population*.

- Non-parametric methods using ranks
 - Ordinal data *and/or...*
 - Central limit theorem does not apply.
- Non-parametric Chi-squared test
 - Can be used for categorical data.

2.1 Single Population

Single Population

- Examine a proportion
 - Parametric: treat data as 0s and 1s, T-test for a single mean.
 - There is a nonparametric test: Binomial distribution.
- Examine the “average” (measure of center) of a single population.
 - Parametric method: T-test for a single mean.
 - Nonparametric methods: Wilcoxon Signed Rank test for a single median.

2.2 Differences in Two Populations

Differences in Two Populations

- Independent Samples
 - Parametric: T-test for difference in means.
 - Nonparametric: Mann-Whitney U-Test - tests whether two populations have the same median.
- Paired samples (Dependent Samples)
 - Parametric: Paired samples T-Test for a difference in means.
 - Nonparametric: Wilcoxon signed rank test for a difference in medians.

2.3 Relationships Between Two Variables

Relationships Between Two Variables

- Parametric method: Pearson linear correlation coefficient.
- Nonparametric method: Spearman correlation.
- Two categorical variables: Chi-squared test of independence.

2.4 Differences in More than Two Populations

Differences in More than Two Populations

- Parametric method: Analysis of Variance (ANOVA)
 - Compares the means of two or more populations.
 - Null hypothesis: all populations have the same mean.
 - Alternative hypothesis: at least one population has a mean different than the others.
- Nonparametric method: Kruskal-Wallis test.
 - Compares the medians of two or more populations.
 - Null hypothesis: all populations have the same median.
 - Alternative hypothesis: at least one population has a median different than the others.

3 Analysis of Variance

3.1 Variance Decomposition

One-Way ANOVA

- Method for testing for significant differences among means from two or more groups.
- Essentially an extension of the t-test for testing the differences between two means.
- Uses measures of *variance* to measure for differences in *means*.
- Total variation in your data is decomposed into two components:
 - **Among-group variation:** variability that is due to differences among groups, also called *explained* variation.
 - **Within-group variation:** total variability within each of the groups, this is unexplained variation.

3.2 Assumptions

Assumptions behind One-way ANOVA F-test

- Randomness: individual observations are assigned to groups *randomly*.
- Independence: individuals in each group are independent from individuals in another group.

- Sufficiently large (?) sample size, or else population must have a normal distribution.
- Homogeneity of variance: the variances of each of the K groups must be equal ($\sigma_1^2 = \sigma_2^2 = \dots \sigma_K^2$).
 - Levene test for homogeneity of variance can be used to test for this.

3.3 Example Using SPSS

Example: Crime Rates

- Data on 47 states from 1960 (I know its old) on the crime rate and a number of factors that may influence the crime rate.
- In particular, I made a variable that put unemployment into categories:
 - Unemployment = 1 if unemployment rate was less than 8%.
 - Unemployment = 2 if unemployment rate was between 8 and 10%.
 - Unemployment = 3 if unemployment rate was greater than 10%.
- I also made a variable that categorized schooling:
 - Schooling = 1 if mean years of schooling for given state was less than 10 years.
 - Schooling = 2 otherwise.
- Is there statistical evidence that the mean crime rate is different among the different categories for the level of unemployment?

FYI: Explanation of all the variables

- R: Crime rate: # of offenses reported to police per million population
- Age: The number of males of age 14-24 per 1000 population
- S: Indicator variable for Southern states (0 = No, 1 = Yes)
- Ed: Mean # of years of schooling x 10 for persons of age 25 or older
- Ex0: 1960 per capita expenditure on police by state and local government
- Ex1: 1959 per capita expenditure on police by state and local government
- LF: Labor force participation rate per 1000 civilian urban males age 14-24
- M: The number of males per 1000 females
- N: State population size in hundred thousands
- NW: The number of non-whites per 1000 population
- U1: Unemployment rate of urban males per 1000 of age 14-24
- U2: Unemployment rate of urban males per 1000 of age 35-39
- W: Median value of transferable goods and assets or family income in tens of \$
- X: The number of families per 1000 earning below 1/2 the median income

Using SPSS to Conduct One-way ANOVA Tests

1. Download and open the dataset `crime.sav` in SPSS.
2. Click on **Analyze** menu, then **Compare Means**, then select **One-Way ANOVA**.
3. Move **Crime rate** to the **Dependent List**.
4. Move **Unemployment** to **Factor**.
5. For extra tests:
 - Click on **Post-hoc** button for tests to compare pair-wise differences in the means.
 - Click on **Options** button for descriptive statistics for for homogeneity of variance test.

One-way ANOVA output

1. **Descriptive Statistics**: shows the mean unemployment rate for each of the three groups, also includes standard deviation, standard error, and confidence intervals. It's nice to present such statistics in your papers.
2. **Levene's Test of Homogeneity of Variances**. The null hypothesis is that the variances are equal.
3. **ANOVA Table**: presents the sum of squares, the mean sum of squares, the F-statistic, and the p-value.
4. **Tukey Tests** for all pairwise comparisons.

4 Kruskal-Wallis Test: Nonparametric Test

4.1 Nonparametric "ANOVA"

Nonparametric One-way ANOVA

- **Kruskal-Wallis Rank Test**: non-parametric technique for testing for differences in the *medians* among two or more groups.
- Like the Mann-Whitney U-test, uses information about the ranks of the observations.
- **Null hypothesis**: All groups have the same median.
- **Alternative hypothesis**: At least one of the medians differs.
- As the sample size gets "large" (over 5 per group some say!), the Kruskal-Wallis test statistic approaches a χ^2 distribution.
- For small sample sizes: possible to compute exact p-values without depending on asymptotic distributions.

4.2 Assumptions

Assumptions for Kruskal-Wallis Test

- Randomness: individual observations are assigned to groups *randomly*.
- Independence: individuals in each group are independent from individuals in another group.
- Only the location (i.e. the center) of the distributions differ among the groups. The populations otherwise have the same distribution.

4.3 Example Using SPSS

Using SPSS to Conduct Kruskal-Wallis Test

1. Click on **Analyze** menu, then **Nonparametric Tests**, then select **K-Independent Samples**.
2. Move **Crime rate** to **Test Variable List**.
3. Move **Unemployment** to **Grouping Variable**.
4. Make sure **Kruskal-Wallis H** text box is selected.
5. Click on **Exact** button if you need exact p-values.
6. Click **OK**!
7. Results show average ranks for each group and χ^2 test statistic and p-values.